

Minimal Length and the Existence of Some Infinitesimal Quantities in Quantum Theory and Gravity

A.E.Shalyt-Margolin ¹

*National Centre of Particles and High Energy Physics, Pervomayskaya
Str. 18, Minsk 220088, Belarus*

PACS: 03.65, 05.20

Keywords: minimal length, infinitesimal quantities, discrete parameters

Abstract

In this work it is demonstrated that, provided a theory involves a minimal length, this theory must be free from such infinitesimal quantities as infinitely small variations in surface of the holographic screen, its volume, and entropy. The corresponding infinitesimal quantities in this case must be replaced by the «minimal variations possible» – finite quantities dependent on the existent energies. As a result, the initial low-energy theory (quantum theory or general relativity) inevitably must be replaced by a minimal-length theory that gives very close results but operates with absolutely other mathematical apparatus.

At the present time all high-energy generalizations (limits) of the basic «components» in fundamental physics (quantum theory [1] and gravity [2]) of necessity lead to a minimal length on the order of the Planck length $l_{min} \propto l_P$. This follows from a string theory [3]– [6], loop quantum gravity [7], and other approaches [8]–[22].

But it is clear that, provided a minimal length exists, it is existent at all the energy scales and not at high (Planck's) scales only.

What is inferred on this basis for real physics? At least, it is suggested that the use of infinitesimal quantities dx_μ in a mathematical apparatus of both quantum theory and gravity is incorrect, despite the fact that both these theories give the results correlating well with the experiment (for example,

¹E-mail: a.shalyt@mail.ru; alexm@hep.by

[23]).

Indeed, in all cases the infinitesimal quantities dx_μ bring about an infinitely small length ds [2]

$$ds^2 = g_{\mu\nu} dx_\mu dx_\nu \quad (1)$$

that is inexistent because of l_{min} .

The same is true for any function Υ dependent only on different parameters L_i whose dimensions of length of the exponents are equal to or greater than 1 $\nu_i \geq 1$

$$\Upsilon \equiv \Upsilon(L_i^{\nu_i}). \quad (2)$$

Obviously, the infinitely small variation $d\Upsilon$ of Υ is senseless as, according to (2), we have

$$d\Upsilon \equiv d\Upsilon(\nu_i L_i^{\nu_i-1} dL_i). \quad (3)$$

But, because of l_{min} , the infinitesimal quantities dL_i make no sense and hence $d\Upsilon$ makes no sense too.

Instead of these infinitesimal quantities it seems reasonable to denote them as «minimal variations possible» Δ_{min} of the quantity L having the dimension of length, i.e. the quantity

$$\Delta_{min} L = l_{min}. \quad (4)$$

And then

$$\Delta_{min} \Upsilon \equiv \Delta_{min} \Upsilon(\nu_i L_i^{\nu_i-1} \Delta_{min} L_i) = \Delta_{min} \Upsilon(\nu_i L_i^{\nu_i-1} l_{min}). \quad (5)$$

However, the «minimal variations possible» of any quantity having the dimensions of length (4) which are equal to $l_{min} \propto l_P$ require, according to the Heisenberg Uncertainty Principle (HUP) [24], maximal momentum $p_{max} \propto P_{Pl}$ and energy $E_{max} \propto E_P$. Here l_P, P_{Pl}, E_P – Planck's length, momentum, and energy, respectively.

But at low energies (far from the Planck energy) there are no such quantities and hence in essence $\Delta_{min} L = l_{min} \propto l_P$ (4) corresponds to the high-energy (Planck's) case only.

For the energies lower than Planck's energy, the «minimal variations possible» $\Delta_{min} L$ of the quantity L having the dimensions of length must be greater than l_{min} and dependent on the present E

$$\Delta_{min} \equiv \Delta_{min,E}, \Delta_{min,E} L > l_{min}. \quad (6)$$

Besides, as we have a minimal length unit l_{min} , it is clear that any quantity having the dimensions of length is «quantized», i.e. its value measured in the units l_{min} equals an integer number and we have

$$L = N_L l_{min}, \quad (7)$$

where N_L —positive integer number.

The problem is, how the «minimal variations possible» $\Delta_{min,E}$ (6) are dependent on the energy or, what is the same, on the scales of the measured lengths?

To solve the above-mentioned problem, **initially** we can use the Space-Time Quantum Fluctuations (STQF) with regard to quantum theory and gravity. The definition (STQF) is closely associated with the notion of «space-time foam». The notion «space-time foam», introduced by J. A. Wheeler about 60 years ago for the description and investigation of physics at Planck's scales (Early Universe) [26],[27], is fairly settled. Despite the fact that in the last decade numerous works have been devoted to physics at Planck's scales within the scope of this notion, for example [28]–[47], by this time still their no clear understanding of the «space-time foam» as it is.

On the other hand, it is undoubtful that a quantum theory of the Early Universe should be a deformation of the well-known quantum theory.

In my works with the colleagues [48]–[57] I has put forward one of the possible approaches to resolution of a quantum theory at Planck's scales on the basis of the density matrix deformation.

In accordance with the modern concepts, the space-time foam [27] notion forms the basis for space-time at Planck's scales (Big Bang). This object is associated with the quantum fluctuations generated by uncertainties in measurements of the fundamental quantities, inducing uncertainties in any distance measurement. A precise description of the space-time foam is still lacking along with an adequate quantum gravity theory. But for the description of quantum fluctuations we have a number of interesting methods (for example, [59],[37]–[47]).

In what follows, we use the terms and symbols from [39]. Then for the fluctuations $\tilde{\delta}l$ of the distance l we have the following estimate:

$$(\tilde{\delta}l)_\gamma \gtrsim l_P^\gamma l^{1-\gamma} = l_P \left(\frac{l}{l_P}\right)^{1-\gamma} = l \left(\frac{l_P}{l}\right)^\gamma = l \lambda_l^\gamma, \quad (8)$$

or that same one

$$|(\tilde{\delta}l)_\gamma|_{min} = \beta l_P^\gamma l^{1-\gamma} = \beta l_P (\frac{l}{l_P})^{1-\gamma} = \beta l \lambda_l^\gamma, \quad (9)$$

where $0 < \gamma \leq 1$, coefficient β is of order 1 and $\lambda_l \equiv l_P/l$.

From (8),(9), we can derive the quantum fluctuations for all the primary characteristics, specifically for the time $(\tilde{\delta}t)_\gamma$, energy $(\tilde{\delta}E)_\gamma$, and metrics $(\tilde{\delta}g_{\mu\nu})_\gamma$. In particular, for $(\tilde{\delta}g_{\mu\nu})_\gamma$ we can use formula (10) in [39]

$$(\tilde{\delta}g_{\mu\nu})_\gamma \gtrsim \lambda^\gamma. \quad (10)$$

Further in the text is assumed that the theory involves a minimal length on the order of Planck's length

$$l_{min} \propto l_P$$

or that is the same

$$l_{min} = \xi l_P, \quad (11)$$

where the coefficient ξ is on the order of unity too.

In this case the origin of the minimal length is not important. For simplicity, we assume that it comes from the Generalized Uncertainty Principle (GUP) that is an extension of HUP for Planck's energies, where gravity must be taken into consideration [3]–[22]:

$$\Delta x \geq \frac{\hbar}{\Delta p} + \alpha' l_P^2 \frac{\Delta p}{\hbar}. \quad (12)$$

Here α' is the model-dependent dimensionless numerical factor.

Inequality (12) leads to the minimal length $l_{min} = \xi l_P = 2\sqrt{\alpha'} l_P$.

Therefore, in this case replacement of Planck's length by the minimal length in all the above formulae is absolutely correct and is used without detriment to the generality [25]

$$l_P \rightarrow l_{min}. \quad (13)$$

Thus, $\lambda_l \equiv l_{min}/l$ and then (8)–(10) upon the replacement of (13) remain unchanged.

As noted in the overview [39], the value $\gamma = 2/3$ derived in [58, 59] is totally

consistent with the Holographic Principle [60]–[63].

The following points of importance should be noted [25]:

1.1) It is clear that **at Planck's scales, i.e. at the minimal length scales**

$$l \rightarrow l_{min} \quad (14)$$

models for different values of the parameter γ are coincident.

1.2) As noted, specifically in (7), **provided some quantity has a minimal measuring unit, values of this quantity are multiples of this unit.**

Naturally, any quantity having a minimal measuring unit is uniformly discrete.

The latter property is not met, in particular, by the energy E .

As $E \sim 1/l$, where l – measurable scale, **the energy E is a discrete but nonuniform quantity**. It is clear that the difference between the adjacent values of E is the less the lower E . In other words, for $l \gg l_{min}$ i. e.

$$E \ll E_P \quad (15)$$

E becomes a practically continuous quantity.

1.3) In fact, the parameter λ_l was introduced earlier in papers [48]–[57] as a deformation parameter on going from the canonical quantum mechanics to the quantum mechanics at Planck's scales (early Universe) that is considered to be the quantum mechanics with the fundamental length (QMFL):

$$0 < \alpha_x = l_{min}^2/x^2 \leq 1/4, \quad (16)$$

where x is the measuring scale, $l_{min} \sim l_p$.

The deformation is understood as an extension of a particular theory by inclusion of one or several additional parameters in such a way that the initial theory appears in the limiting transition [64]. Obviously, everywhere, apart from the limiting point $\lambda_x = 1$ or $x = l_{min}$, we have

$$\lambda_x = \sqrt{\alpha_x}, \quad (17)$$

From (16) it is seen that at the limiting point $x = l_{min}$ the parameter α_x is not defined due to the appearance of singularity [48]–[57]. But at this point its definition may be extended (regularized).

The parameter α_l has the following clear physical meaning:

$$\alpha_l^{-1} \sim S^{BH}, \quad (18)$$

where

$$S^{BH} = \frac{A}{4l_p^2} \quad (19)$$

is the well-known Bekenstein-Hawking formula for the black hole entropy in the semiclassical approximation [65],[66] for the black-hole event horizon surface A , with the characteristics linear dimension («radius») $R = l$. This is especially obvious in the spherically-symmetric case.

In what follows we use both parameters: λ_x and α_x .

Turning back to the introductory section of this work and to the definition $\Delta_{min,E}L$, we assume the following:

$$|\Delta_{min,E}L| = |(\tilde{\delta}L)_\gamma|_{min}, \quad (20)$$

where $|(\tilde{\delta}L)_\gamma|_{min}$ is from formula (9), γ – fixed parameter from formulae (8), (9), and $E = c\hbar/L$.

In physics, and in thermodynamics in particular, the *extensive quantities or parameters* are those proportional to the mass of a system or to its volume. Proceeding from the definition (2) of the function $\Upsilon(L_i^{\nu_i})$, one can generalize this notion, taking as a **Generalized Extensive Quantity (GEQ)** of some spatial system Ω the function dependent only on the linear dimensions of this system, with the exponents no less than 1.

The function $\Upsilon(L_i^{\nu_i})$, $\nu_i \geq 1$ (2) is **GEQ** of the system Ω with the characteristic linear dimensions L_i ; $i = 1, \dots, n$ or, identically, a sum of the systems Ω_i ; $i = 1, \dots, n$, each of which has its individual characteristic linear dimension L_i .

Then from the initial formulae (2)–(6) it directly follows that, provided the minimal length l_{min} is existent, **there are no** infinitesimal variations of GEQ.

In the first place, this is true for such simplest objects as the n -dimensional

sphere $B_n, n \geq 2$, whose surface area (area of the corresponding hypersphere S_n) and volume V_n represent GEQs and are equal to the following:

$$S_n = nC_n R^{n-1}; V_n = C_n R^n, \quad (21)$$

where R – radius of a sphere the length of which is a characteristic linear size, $C_n = \pi^{n/2}/\Gamma(\frac{n}{2} + 1)$, and $\Gamma(x)$ is a gamma-function.

Of course, the same is true for the n -dimensional cube (or hypercube) A_n ; its surface area and its volume are GEQs, and a length of its edge is a characteristic linear dimension.

Provided l_{min} exists, there are no infinitesimal increments for both the surface area and volume of A_n or B_n ; only minimal variations possible for these quantities are the case.

In what follows we consider only the spatial systems whose surface areas and volumes are GEQs.

Let us consider a simple but very important example of gravity in horizon spaces.

Gravity and thermodynamics of horizon spaces and their interrelations are currently most actively studied [67]–[79]. Let us consider a relatively simple illustration – the case of a static spherically-symmetric horizon in space-time, the horizon being described by the metric

$$ds^2 = -f(r)c^2 dt^2 + f^{-1}(r)dr^2 + r^2 d\Omega^2. \quad (22)$$

The horizon location will be given by a simple zero of the function $f(r)$, at the radius $r = a$.

This case is studied in detail by T.Padmanabhan in his works [67, 78] and by the author of this paper in [80]. We use the notation system of [78]. Let, for simplicity, the space be denoted as $\mathcal{H}$.

It is known that for horizon spaces one can introduce the temperature that can be identified with an analytic continuation to imaginary time. In the case under consideration ([78], eq.(116))

$$k_B T = \frac{\hbar c f'(a)}{4\pi}. \quad (23)$$

Therewith, the condition $f(a) = 0$ and $f'(a) \neq 0$ must be fulfilled. Then at the horizon $r = a$ Einstein's equations have the form

$$\frac{c^4}{G} \left[\frac{1}{2} f'(a) a - \frac{1}{2} \right] = 4\pi P a^2 \quad (24)$$

where $P = T_r^r$ is the trace of the momentum-energy tensor and radial pressure.

Now we proceed to the variables $\ll \alpha \gg$ from the formula (16) to consider (24) in a new notation, expressing a in terms of the corresponding deformation parameter α . In what follows we omit the subscript in formula (16) of α_x , where the context implies which index is the case. In particular, here we use α instead of α_a . Then we have

$$a = l_{min} \alpha^{-1/2}. \quad (25)$$

Therefore,

$$f'(a) = -2l_{min}^{-1} \alpha^{3/2} f'(\alpha). \quad (26)$$

Substituting this into (24) we obtain in the considered case of Einstein's equations in the $\ll \alpha$ -representation $\gg$ the following [80]:

$$\frac{c^4}{G} \left(-\alpha f'(\alpha) - \frac{1}{2} \right) = 4\pi P \alpha^{-1} l_{min}^2. \quad (27)$$

Multiplying the left- and right-hand sides of the last equation by α , we get

$$\frac{c^4}{G} \left(-f'(\alpha) \alpha^2 - \frac{1}{2} \alpha \right) = 4\pi P l_{min}^2. \quad (28)$$

L.h.s. of (28) is dependent on α . Because of this, r.h.s. of (28) must be dependent on α as well, i. e. $P = P(\alpha)$, i.e

$$\frac{c^4}{G} \left(-f'(\alpha) \alpha^2 - \frac{1}{2} \alpha \right) = 4\pi P(\alpha) l_{min}^2. \quad (29)$$

Note that in this specific case the parameter α within constant factors is coincident with the Gaussian curvature K_a [81] corresponding to a :

$$\frac{l_{min}^2}{a^2} = l_{min}^2 K_a. \quad (30)$$

Substituting r.h.s of (30) into (29), we obtain the Einstein equation on horizon, in this case in terms of the Gaussian curvature

$$\frac{c^4}{G}(-f'(K_a)K_a^2 - \frac{1}{2}K_a) = 4\pi P(K_a). \quad (31)$$

This means that up to the constants

$$-f'(K_a)K_a^2 - \frac{1}{2}K_a = P(K_a), \quad (32)$$

i.e. the Gaussian curvature K_a is a solution of Einstein equations in this case. Then we examine different cases of the solution (32) with due regard for considerations of formula (20).

2.1) First, let us assume that $a \gg l_{min}$. As, according (7), the radius a is quantized, we have $a = N_a l_{min}$ with the natural number $N_a \gg 1$. Then it is clear that the Gaussian curvature $K_a = 1/a^2 \approx 0$ takes a (nonuniform) discrete series of values close to zero, and, within the factor $1/l_{min}^2$, this series represents inverse squares of natural numbers

$$(K_a) = (\frac{1}{N_a^2}, \frac{1}{(N_a \pm 1)^2}, \frac{1}{(N_a \pm 2)^2}, \dots). \quad (33)$$

Let us return to formulas (9),(20) for $l = a$

$$|((\tilde{\delta}a)_\gamma)_{min}| = \beta N_a l_{min} N_a^{-\gamma} = \beta N_a^{1-\gamma} l_{min}, \quad (34)$$

where β in this case contains the proportionality factor that relates l_{min} and l_P .

Then, according to (20), $a_{\pm 1}$ is a measurable value of the radius r following after a , and we have

$$(a_{\pm 1})_\gamma \equiv a \pm ((\tilde{\delta}a)_\gamma)_{min} = a \pm \beta N_a^{1-\gamma} l_{min} = N_a(1 \pm \beta N_a^{-\gamma}) l_{min}. \quad (35)$$

But, as $N_a \gg 1$, for sufficiently large N_a and fixed γ , the bracketed expression in r.h.s. (35) is close to 1:

$$1 \pm \beta N_a^{-\gamma} \approx 1. \quad (36)$$

Obviously, we get

$$\lim_{N_a \rightarrow \infty} (1 \pm \beta N_a^{-\gamma}) \rightarrow 1. \quad (37)$$

As a result, the Gaussian curvature $K_{a_{\pm 1}}$ corresponding to $r = a_{\pm 1}$

$$K_{a_{\pm 1}} = 1/a_{\pm 1}^2 \propto \frac{1}{N_a^2 (1 \pm \beta N_a^{-\gamma})^2} \quad (38)$$

in the case under study is only slightly different from K_a .

And this is the case for sufficiently large values of N_a , for any value of the parameter γ , for $\gamma = 1$ as well, corresponding to the absolute minimum of fluctuations $\approx l_{min}$, or more precisely – to βl_{min} . However, as all the quantities of the length dimension are quantized and the factor β is on the order of 1, actually we have $\beta = 1$.

Because of this, provided the minimal length is involved, l_{min} (9) is read as

$$|(\tilde{\delta}l)_1|_{min} = l_{min}. \quad (39)$$

But, according to (11), $l_{min} = \xi l_P$ is on the order of Planck's length, and it is clear that the fluctuation $|(\tilde{\delta}l)_1|_{min}$ corresponds to Planck's energies and Planck's scales. The Gaussian curvature K_a , due to its smallness ($K_a \ll 1$ up to the constant factor l_{min}^{-2}) and smooth variations independent of γ (formulas (35)–(38)), is **insensitive** to the differences between various values of γ .

Consequently, for sufficiently small Gaussian curvature K_a we can take any parameter from the interval $0 < \gamma \leq 1$ as γ .

It is obvious that the case $\gamma = 1$, i. e. $|(\tilde{\delta}l)_1|_{min} = l_{min}$, is associated with infinitely small variations da of the radius $r = a$ in the Riemannian geometry.

Since then K_a is varying practically continuously, in terms of K_a up to the constant factor we can obtain the following expression:

$$d[L(K_a)] = d[P(K_a)], \quad (40)$$

Where have

$$L(K_a) = -f'(K_a)K_a^2 - \frac{1}{2}K_a, \quad (41)$$

i. e. l.h.s of (31) (or (32)).

But in fact, as in this case the energies are low, it is more correct to consider

$$L((K_{a\pm 1})_\gamma) - L(K_a) = [P(K_{a\pm 1})_\gamma] - [P(K_a)] \equiv F_\gamma[P(K_a)], \quad (42)$$

where $\gamma < 1$, rather than (40).

In view of the foregoing arguments 2.1), the difference between (42) and (40) is insignificant and it is perfectly correct to use (40) instead of (42).

2.2) Now we consider the opposite case or the transition to the **ultraviolet limit**

$$a \rightarrow l_{min} = \kappa l_{min}, \quad (43)$$

i.e.

$$a = \kappa l_{min}. \quad (44)$$

Here κ is on the order of 1.

Taking into consideration point 1.1) stating that in this case models for different values of the parameter γ are coincident, by formula (39) for any γ we have

$$|(\tilde{\delta}l)_\gamma|_{min} = |(\tilde{\delta}l)_1|_{min} = l_{min}. \quad (45)$$

But in this case the Gaussian curvature K_a is not a «small value» continuously dependent on a , taking, according to (38), a discrete series of values $K_a, K_{a\pm 1}, K_{a\pm 2}, \dots$

Yet (24), similar to (31) ((32)), is valid in the semiclassical approximation only, i.e. at **low energies**.

Then in accordance with the above arguments, the limiting transition to **high energies** (43) gives a discrete chain of equations or a single equation with a discrete set of solutions as follows:

$$-f'(K_a)K_a^2 - \frac{1}{2}K_a = \Theta(K_a);$$

$$-f'(K_{a\pm 1})K_{a\pm 1}^2 - \frac{1}{2}K_{a\pm 1} = \Theta(K_{a\pm 1});$$

and so on. Here $\Theta(K_a)$ – some function that in the limiting transition to low

energies must reproduce the low-energy result to a high degree of accuracy, i.e. $P(K_a)$ appears for $a \gg l_{min}$ from formula (32)

$$\lim_{K_a \rightarrow 0} \Theta(K_a) = P(K_a). \quad (46)$$

In general, $\Theta(K_a)$ may lack coincidence with the high-energy limit of the momentum-energy tensor trace (if any):

$$\lim_{a \rightarrow l_{min}} P(K_a). \quad (47)$$

At the same time, when we naturally assume that the Static Spherically-Symmetric Horizon Space-Time with the radius of several Planck's units (44) is nothing else but a micro black hole, then the high-energy limit (47) is existing and the replacement of $\Theta(K_a)$ by $P(K_a)$ in r.h.s. of the foregoing equations is possible to give a hypothetical gravitational equation for the event horizon micro black hole. But a question arises, for which values of the parameter a (44) (or K_a) this is valid and what is a minimal value of the parameter $\gamma = \gamma(a)$ in this case.

In all the cases under study, 2.1) and 2.2), the deformation parameter α_a (16) (λ_a (17)) is, within the constant factor, coincident with the Gaussian curvature K_a (respectively $\sqrt{K_a}$) that is in essence continuous in the low-energy case and discrete in the high-energy case.

In this way the above-mentioned example shows that, despite the absence of infinitesimal spatial-temporal increments owing to the existence of l_{min} and the essential «discreteness» of a theory, this discreteness at low energies is not «felt», the theory being actually continuous. The indicated discreteness is significant only in the case of high (Planck's) energies.

In [78] it is shown that the Einstein Equation for horizon spaces in the differential form may be written as a thermodynamic identity (the first principle of thermodynamics) ([78], formula (119)):

$$\underbrace{\frac{\hbar c f'(a)}{4\pi}}_{k_B T} \underbrace{\frac{c^3}{G \hbar} d \left(\frac{1}{4} 4\pi a^2 \right)}_{dS} - \underbrace{\frac{1}{2} \frac{c^4 da}{G}}_{-dE} = \underbrace{P d \left(\frac{4\pi}{3} a^3 \right)}_{P dV}. \quad (48)$$

where, as noted above, T – temperature of the horizon surface, S – corresponding entropy, E – internal energy, V – space volume. Note that, because of the existing l_{min} , practically all quantities in (48) (except of T) represent GEQ. Apparently, the radius of a sphere $r = a$, its volume V , and entropy represent such quantities:

$$S = \frac{4\pi a^2}{4l_P^2} = \frac{\pi a^2}{l_P^2}, \quad (49)$$

within the constant factor $1/4l_P^2$ equal to a sphere with the radius a . Because of this, there are no infinitesimal increments of these quantities, i.e. da, dV, dS . And, provided l_{min} is involved, the Einstein equation for the above-mentioned case in the differential form (48) makes no sense and is useless. If da may be, purely formally, replaced by l_{min} , then, as the quantity l_{min} is fixed, it is obvious that $\langle dS \rangle$ and $\langle dV \rangle$ in (48) will be growing as a and a^2 , respectively. And at low energies, i.e. for large values of $a \gg l_{min}$, this naturally leads to infinitely large rather than infinitesimal values.

In a similar way it is easily seen that the «Entropic Approach to Gravity» [82] in the present formalism is invalid within the scope of the minimal length theory. In fact, the «main instrument» in [82] is a formula for the infinitesimal variation dN in the bit numbers N on the holographic screen $\mathcal{S}$ with the radius R and with the surface area A ([82], formula (4.18)):

$$dN = \frac{c^3}{G\hbar} dA = \frac{dA}{l_P^2}. \quad (50)$$

As $N = A/l_P^2$, and A represents GEQ, it is clear that N is also GEQ and hence neither dA nor dN makes sense.

It is obvious that infinitesimal variations of the screen surface area dA are possible only in a continuous theory involving no l_{min} .

When $l_{min} \propto l_P$ is involved, the minimal variation ΔA is evidently associated with a minimal variation in the radius R

$$R \rightarrow R \pm l_{min} \quad (51)$$

is dependent on R and growing as R for $R \gg l_{min}$

$$\Delta_{\pm} A(R) = (A(R \pm l_{min}) - A(R)) \propto \left(\frac{\pm 2R}{l_{min}} + 1 \right) = \pm 2N_R + 1, \quad (52)$$

where $N_R = R/l_{min}$, as indicated above.

But, as noted above, a minimal increment of the radius R equal to $|\Delta_{min}R| = l_{min} \propto l_P$ corresponds only to the case of maximal (Planck's) energies or, what is the same, to the parameter $\gamma = 1$ in formula (20). However, in [82] the considered low energies are far from the Planck energies and hence in this case in (20) $\gamma < 1$, (51), and (52) are respectively replaced by

$$R \rightarrow R \pm N_R^{1-\gamma} l_{min} \quad (53)$$

and

$$\Delta_{\pm} A(R) = (A(R \pm N_R^{1-\gamma} l_{min}) - A(R)) \propto \pm N_R^{2-\gamma} + N_R^{2-2\gamma} = N_R^{2-2\gamma} (\pm N_R^{\gamma} + 1). \quad (54)$$

An increase of r.h.s in (54) with the growth of R (or identically of N_R) for $R \gg l_{min}$ is obvious.

So, if l_{min} is involved, formula (4.18) from [82] makes no sense similar to other formulae derived on its basis (4.19), (4.20), (4.22), (5.32)–(5.34), ... in [82] and similar to the derivation method for Einstein's equations proposed in this work.

Proceeding from the principal parameters of this work α_l (or λ_l), the fact is obvious and is supported by the formula (18) given in this paper, meaning that

$$\alpha_R^{-1} \sim A, \quad (55)$$

i.e. small variations of α_R (low energies) result in large variations of α_R^{-1} , as indicated by formula (52).

In fact, we have a **no-go theorems**.

The last statements concerning dS, dN may be explicitly interpreted using the language of a quantum information theory as follows:

due to the existence of the minimal length l_{min} , the minimal area l_{min}^2 and volume l_{min}^3 are also involved, and that means «quantization» of the areas and volumes. As, up to the known constants, the «bit number» N from (50) and the entropy S from (49) are nothing else but

$$S = \frac{A}{4l_{min}^2}, N = \frac{A}{l_{min}^2}, \quad (56)$$

it is obvious that there is a «minimal measure» for the «amount of data» that may be referred to as «one bit» (or «one qubit»).

The statement that there is no such quantity as dN (and respectively dS) is equivalent to claiming the absence of 0.25 bit, 0.001 bit, and so on.

This inference completely conforms to the Hooft-Susskind Holographic Principle (HP) [60]–[63] that includes two main statements:

- (a) All information contained in a particular spatial domain is concentrated at the boundary of this domain.
- (b) A theory for the boundary of the spatial domain under study should contain maximally one degree of freedom per Planck's area l_P^2 .

In fact (but not explicitly) HP implicates the existence of $l_{min} = l_P$. The existence of $l_{min} \propto l_P$ totally conforms to HP, providing its generalization. Specifically, without the loss of generality, l_P^2 in point (b) may be replaced by l_{min}^2 .

So, the principal inference of this work is as follows:

provided the minimal length l_{min} is involved, its existence must be taken into consideration not only at high but also at low energies, both in a quantum theory and in gravity. This becomes apparent by rejection of the infinitesimal quantities associated with the spatial-temporal variations $dx_\mu, \dots$. In other words, with the involvement of l_{min} , the General Relativity (GR) must be replaced by a (still unframed) minimal-length gravitation theory that may be denoted as $Grav^{l_{min}}$. In their results GR and $Grav^{l_{min}}$ should be very close but, as regards their mathematical apparatus (instruments), these theories are absolutely different.

Besides, $Grav^{l_{min}}$ should offer a rather natural transition from high to low energies

$$[N_L \approx 1] \rightarrow [N_L \gg 1] \quad (57)$$

and vice versa

$$[N_L \gg 1] \rightarrow [N_L \approx 1], \quad (58)$$

where N_L – integer from formula (7) determining the characteristics scale of the lengths L (energies $E \sim 1/L \propto 1/N_L$).

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this article.

References

- [1] M.E. Peskin, D.V. Schroeder, *An Introduction to Quantum Field Theory*, Addison-Wesley Publishing Company, 1995.
- [2] R. M. Wald, *General Relativity*, University of Chicago Press, Chicago, Ill, USA, 1984.
- [3] G. A. Veneziano, “Stringy nature needs just two constants,” *Europhysics Letters*, vol. 2, no. 3, pp. 199–211, 1986.
- [4] D. Amati, M. Ciafaloni, and G. Veneziano, “Can spacetime be probed below the string size?” *Physics Letters B*, vol. 216, no. 1-2, pp. 41–47, 1989.
- [5] E. Witten, “Reflections on the fate of spacetime,” *Physics Today*, vol. 49, no. 4, pp. 24–28, 1996.
- [6] Joseph Polchinski, *String Theory*, Vol. I, II, Cambridge University Press, 1998.
- [7] Carlo Rovelli, “Loop quantum gravity: the first twenty five years,” *Classical and Quantum Gravity*, vol. 28, p.p. 153002, 2011.
- [8] G. Amelino-Camelia, “Quantum Spacetime Phenomenology,” *Living Rev. Rel.*, vol. 16, pp. 5–129, 2013; arxiv.org/abs/gr-qc/0806.0339.
- [9] L. Garay, “Quantum gravity and minimum length,” *International Journal of Modern Physics A*, vol. 10, no. 2, pp. 145–146, 1995.
- [10] G. Amelino-Camelia, L. Smolin, “Prospects for constraining quantum gravity dispersion with near term observations,” *Phys. Rev. D*, vol. 80, 084017, 2009.

- [11] G.Gubitosi et al., "A constraint on planck-scale modifications to electrodynamics with CMB polarization data," *JCAP*, vol. 0908, pp.021-034,2009.
- [12] G.Amelino-Camelia," Building a case for a planck-scale-deformed boost action: The planck-scale particle-localization limit." *Int.J.Mod.Phys.D.* vol.14,pp.2167–2180,2005.
- [13] S. Hossenfelder et al., "Signatures in the Planck Regime," *Phys. Lett.B*, vol.575,pp.85–99,2003.
- [14] S. Hossenfelder,"Running Coupling with Minimal Length," *Phys.Rev.D*,vol.70,p.105003,2004;
- [15] S. Hossenfelder,"Self-consistency in Theories with a Minimal Length" *Class. Quant. Grav.*, vol.23,pp.1815–1821,2006.
- [16] S. Hossenfelder,"Minimal Length Scale Scenarios for Quantum Gravity," *Living Rev.Rel.*, vol.16,pp.2–91,2013.
- [17] R. J. Adler and D. I. Santiago, "On gravity and the uncertainty principle," *Modern Physics Letters A*, vol. 14, no. 20, pp. 1371– 1378, 1999.
- [18] D. V. Ahluwalia, "Wave-particle duality at the Planck scale: freezing of neutrino oscillations," *Physics Letters A.*, vol. 275, no. 1-2,pp.31–35,2000.
- [19] D. V. Ahluwalia, "Interface of gravitational and quantum realms," *Modern Physics Letters A*, vol. 17, no. 15–17,pp.1135–1145,2002.
- [20] M.Maggiore, "The algebraic structure of the generalized uncertainty principle," *Physics Letters B*, vol. 319, no. 1–3, pp. 83–86, 1993.
- [21] A. Kempf, G. Mangano, and R. B. Mann, "Hilbert space representation of the minimal length uncertainty relation," *Physical Review D*, vol. 52, no. 2, pp. 1108–1118, 1995.
- [22] Kouros Nozari, Amir Etemadi,"Minimal length, maximal momentum and Hilbert space representation of quantum mechanics," *Physical Review D*,vol.85,p.104029,2012.

- [23] R. Penrose, *Quantum Theory and Space-Time*, Fourth Lecture in *Stephen Hawking and Roger Penrose, The Nature of Space and Time*, Princeton University Press, 1996.
- [24] W. Heisenberg, "Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik," *Zeitschrift für Physik.*, vol.43, no 3-4, pp.172–198, 1927; English translation: J. A. Wheeler and H. Zurek, *Quantum Theory and Measurement* Princeton Univ. Press, 1983, pp. 62-84.
- [25] A.E. Shalyt-Margolin, "Space-Time Fluctuations, Quantum Field Theory with UV-cutoff and Einstein Equations," *Nonlinear Phenomena in Complex Systems*, vol.17, no 2, pp.138–146, 2014.
- [26] J. A. Wheeler, *Geometrodynamics*, Academic Press, New York and London, 1962.
- [27] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* Freeman, San Francisco, 1973.
- [28] Remo Garattini, "A Spacetime Foam approach to the cosmological constant and entropy," *Int. J. Mod. Phys. D.*, vol.4, pp.635–652, 2002.
- [29] Remo Garattini, "A Spacetime Foam Approach to the Schwarzschild-de Sitter Entropy," *Entropy*, vol.2, pp.26–38, 2000.
- [30] Remo Garattini, "Entropy and the cosmological constant: a spacetime-foam approach," *Nucl.Phys.Proc.Suppl.*, vol.88, pp.297–300, 2000.
- [31] Remo Garattini, "Entropy from the foam," *Phys.Lett.B.*, vol.459, pp.461–467, 1999.
- [32] Fabio Scardigli, "Black Hole Entropy: a spacetime foam approach," *Class.Quant.Grav.*, vol.14, pp.1781–1793, 1997.
- [33] Fabio Scardigli, "Generalized Uncertainty Principle in Quantum Gravity from Micro-Black Hole Gedanken Experiment," *Phys.Lett.B.*, vol.452, pp.39–44, 1999.

- [34] Fabio Scardigli,"Gravity coupling from micro-black holes," *Nucl.Phys.Proc.Suppl.*,vol.88,pp.291–294 2000.
- [35] Luis J. Garay,"Thermal properties of spacetime foam," *Phys.Rev.D.*,vol.58, p.124015,1998.
- [36] Luis J. Garay,"Spacetime foam as a quantum thermal bath," *Phys.Rev.Lett.*, vol.80,2508–2511,1998.
- [37] Y. J. Ng, H. van Dam,"Measuring the foaminess of space-time with gravity-wave interferometers," *Found. Phys.*,vol.30,pp.795–805,2000.
- [38] Y. J. Ng,"Spacetime foam," *Int. J. Mod. Phys. D.*, vol.11,1585–1590,2002.
- [39] Y. J. Ng,"Selected topics in Planck-scale physics," *Mod.Phys.Lett.A.*, vol.18,pp.1073–1098,2003.
- [40] Y. J. Ng,"Quantum Foam," arxiv.org/abs/gr-qc/0401015.
- [41] Y. J. Ng, H. van Dam,"Spacetime Foam, Holographic Principle, and Black Hole Quantum Computers," *Int.J.Mod.Phys.A*.vol.20,pp.1328–1335,2005.
- [42] W.A. Christiansen, Y. Jack Ng and H. van Dam,"Probing spacetime foam with extragalactic sources," *Phys.Rev.Lett.*, vol.96, p.051301,2006.
- [43] Y. Jack Ng,"Holographic Foam, Dark Energy and Infinite Statistics," *Phys.Lett.B.*,vol.657,10–14,2007.
- [44] Y. Jack Ng,"Spacetime Foam and Dark Energy," *AIP Conf.Proc.*,vol.1115,pp.74–79,2009.
- [45] A. Wayne Christiansen, David J. E. Floyd, Y. Jack Ng, Eric S. Perlman,"Limits on Spacetime Foam," *Phys.Rev.D.*, vol.83,p.084003,2011.
- [46] G. Amelino-Camelia,"An interferometric gravitational wave detector as a quantum-gravity apparatus," *Nature*,vol.398,pp. 216–218,1999.

- [47] L. Diosi and B. Lukacs, "On the minimum uncertainties of space-time geodesics", *Phys. Lett. A* **142**, pp.331–334,1989.
- [48] A. E. Shalyt-Margolin and J. G. Suarez, "Quantum mechanics of the early universe and its limiting transition," <http://arxiv.org/abs/gr-qc/0302119>.
- [49] A. E. Shalyt-Margolin and J. G. Suarez, "Quantum mechanics at Planck's scale and density matrix," *International Journal of Modern Physics D*, vol. 12, no. 7, pp. 1265–1278, 2003.
- [50] A. E. Shalyt-Margolin and A. Y. Tregubovich, "Deformed density matrix and generalized uncertainty relation in thermodynamics," *Modern Physics Letters A*, vol. 19, no. 1, pp. 71–82, 2004.
- [51] A. E. Shalyt-Margolin, "Non-unitary and unitary transitions in generalized quantum mechanics, new small parameter and information problemsolving," *Modern Physics Letters A*, vol. 19, no. 5, pp. 391–403, 2004.
- [52] A. E. Shalyt-Margolin, "Pure states, mixed states and Hawking problem in generalized quantum mechanics," *Modern Physics Letters A*, vol. 19, no. 27, pp. 2037–2045, 2004.
- [53] A. E. Shalyt-Margolin, "The universe as a nonuniform lattice in finite-volume hypercube: I. Fundamental definitions and particular features," *International Journal of Modern Physics D*, vol. 13, no. 5, pp. 853–864, 2004.
- [54] A. E. Shalyt-Margolin, "The Universe as a nonuniformlattice in the finite-dimensional hypercube. II. Simple cases of symmetry breakdown and restoration," *International Journal of Modern Physics A*, vol. 20, no. 20-21, pp. 4951–4964, 2005.
- [55] A. E. Shalyt-Margolin and V. I. Strazhev, "The density matrix deformation in quantum and statistical mechanics of the early universe," in *Proceedings of the 6th International Symposium "Frontiers of Fundamental and Computational Physics"*, B. G. Sidharth, Ed., pp. 131–134, Springer, Berlin, Germany, 2006

- [56] A. Shalyt-Margolin, “Entropy in the present and early universe: new small parameters and dark energy problem,” *Entropy*, vol. 12, no. 4, pp. 932–952, 2010.
- [57] A. E. Shalyt-Margolin, “The density matrix deformation in physics of the early universe and some of its implications,” in *Quantum Cosmology Research Trends*, A. Reimer, Ed., Horizons in World Physics no. 246, pp. 49–92, Nova Science, Hauppauge, NY, USA, 2005.
- [58] E.P. Wigner, “Relativistic Invariance and Quantum Phenomena,” *Rev. Mod. Phys.*, vol. 29, pp. 255–281, 1957.
- [59] H. Salecker and E.P. Wigner, “Quantum Limitations of the Measurement of Space-Time Distances,” *Phys. Rev.*, vol. 109, pp. 571–584, 1958.
- [60] G. ’t Hooft, “Dimensional reduction in quantum gravity,” <http://arxiv.org/abs/gr-qc/9310026>.
- [61] G. ’t Hooft, “The holographic principle,” <http://arxiv.org/abs/hep-th/0003004>.
- [62] L. Susskind, “The world as a hologram,” *Journal of Mathematical Physics*, vol. 36, no. 11, pp. 6377–6396, 1995.
- [63] R. Bousso, “The holographic principle,” *Reviews of Modern Physics*, vol. 74, no. 3, pp. 825–874, 2002.
- [64] L. Faddeev, “Mathematical view of the evolution of physics,” *Priroda*, vol. 5, pp. 11–16, 1989.
- [65] J. D. Bekenstein, “Black holes and entropy,” *Physical Review D.*, vol. 7, pp. 2333–2346, 1973.
- [66] S. W. Hawking, “Black holes and thermodynamics,” *Physical Review D*, vol. 13, no. 2, pp. 191–197, 1976.
- [67] T. Padmanabhan, “Classical and Quantum Thermodynamics of horizons in spherically symmetric spacetimes,” *Class. Quant. Grav.*, vol. 19, pp. 5387–5408, 2002.

- [68] T. Padmanabhan, “A new perspective on gravity and dynamics of space-time,” *International Journal of Modern Physics D*, vol. 14, no. 12, pp. 2263–2270, 2005.
- [69] T. Padmanabhan, “The holography of gravity encoded in a relation between entropy, horizon area, and action for gravity,” *General Relativity and Gravitation*, vol. 34, no. 12, pp. 2029–2035, 2002.
- [70] T. Padmanabhan, “Holographic gravity and the surface term in the Einstein-Hilbert action,” *Brazilian Journal of Physics.*, vol.35,no.2,pp.362–372,2005.
- [71] T. Padmanabhan, “Gravity: a new holographic perspective,” *International Journal of Modern Physics D*, vol. 15, no. 10, pp. 1659–1676, 2006.
- [72] A. Mukhopadhyay and T. Padmanabhan, “Holography of gravitational action functionals,” *Physical Review D*, vol. 74, no. 12, Article ID 124023, 15 pages, 2006.
- [73] T. Padmanabhan, “Dark energy and gravity,” *General Relativity and Gravitation*, vol. 40, no. 2-3, pp. 529–564, 2008.
- [74] T. Padmanabhan and A. Paranjape, “Entropy of null surfaces and dynamics of spacetime,” *Physical Review D*, vol. 75, no. 6, Article ID 064004, 16 pages, 2007.
- [75] T. Padmanabhan, “Gravity as an emergent phenomenon: a conceptual description,” *AIP Conference Proceedings*, vol. 939, pp. 114–123, 2007.
- [76] T. Padmanabhan, “Gravity and the thermodynamics of horizons,” *Physics Reports*, vol. 406, no. 2, pp. 49–125, 2005.
- [77] A. Paranjape, S. Sarkar, and T. Padmanabhan, “Thermodynamic route to field equations in Lanczos-Lovelock gravity,” *Physical Review D*, vol. 74, no. 10, Article ID 104015, 9 pages, 2006.
- [78] T. Padmanabhan, “Thermodynamical aspects of gravity: new insights,” *Reports on Progress in Physics*, vol. 73, Article ID 046901, 2010.

- [79] T. Padmanabhan, “Equipartition of energy in the horizon degrees of freedom and the emergence of gravity,” *Modern Physics Letters A*, vol. 25, no. 14, pp. 1129–1136, 2010.
- [80] A. E. Shalyt-Margolin, “Quantum theory at planck scale, limiting values, deformed gravity and dark energy problem,” *International Journal of Modern Physics D*, vol. 21, no. 2, Article ID 1250013, 20 pages, 2012.
- [81] S. Kobayashi, K. Nomozu, *Foundations of Differential Geometry*, V.II, Interscience Publishers, New York-London-Sydney 1969.
- [82] E. Verlinde, “On the origin of gravity and the laws of Newton,” *Journal of High Energy Physics*, vol. 2011, no. 4, article 29, 2011.